



ARTHROSCOPIC CAPSULAR RELEASE (MUA) REHABILITATION GUIDELINES

INTRODUCTION

The physiotherapy programme will need to be **individualised** for each patient, all exercises should be performed without pain and the details of specific restrictions will be in the post-operative instructions. If you have not received these please ring the consultant's secretary.

Emphasise to the patient the importance allowing soft-tissue healing in the first two phases. Alongside this we aim to regain passive range of movement (PROM) and good scapula-humeral rhythm (SHR) before moving onto strengthening in stage 3. The milestones may be used to assess whether you feel the patient is making good progress or not.

Shoulder rehabilitation is more than strength-training of the shoulder muscles alone. The (neuromuscular) rehabilitation addresses the whole shoulder girdle, upper extremity, core stability and training of the kinetic chain.

POST SURGERY

Phase I (1 – 14 days)

Goals:

- Management of pain, inflammation and muscle inhibition
 - Analgesics, NSAID's, ice, sling, passive movement and posture
- Teach shoulder girdle control/setting and relaxation
 - Retraction and depression
- Increase PROM **as tolerated**
 - Pendulum with good SHR
 - ER/IR
- Progress to active range of movement (AROM) **as tolerated/not into pain** and ensuring good SHR
- Hand, wrist, elbow and neck range of movement (ROM) exercises as required
- Advice on sleeping position
- Return to light, non-repetitive functional activities
- Prevent muscle atrophy
 - Sub-maximal, pain-free isometrics in neutral (<30% maximal voluntary contraction) as tolerated

Precautions:

- Sling usually for 1-2 days for comfort as required
- Keep wounds clean and dry

Milestones:

- Pain, inflammation and muscle inhibition well managed
- Return to pre-operative sleep patterns
- Good scapula setting
- 75% of intra-operative, post-MUA/capsular release ROM

Arthroscopic Capsular Release post-op Protocol		Classification: Company Confidential
Document type: Post-op Protocol	Page: 1 of 3	Author: Darren James
Effective Date: March 2022	Version: 2.1	Next Review: March 2024

- **Phase 2 (15 Days – 6 weeks)**

Goals:

- Continue to manage and reduce pain, inflammation and muscle inhibition
 - As phase 1
 - Alternate treatment strategies as appropriate e.g. manual therapy techniques, taping
- Once stitches have been removed begin scar massage
- Restore intra-operative, post-MUA/capsular release PROM and AROM **as tolerated**
- Re-establish dynamic shoulder stability
 - Ensure good SHR through PROM and use it to guide progression of AROM
- Progress to maximal, pain-free isometrics
- Gradually introduce proprioceptive exercise, lower limb and core strengthening as required and light, early-stage sport-specific exercises
- Gradually return to light, non-repetitive functional activities

Milestones:

- Minimal pain, inflammation and muscle inhibition
- 100% of intra-operative, post-MUA/capsular release PROM
- Good SHR with AROM

Phase 3 (7 - 14 weeks)

Goals:

- Full PROM
 - Introduce multi-directional stretching into end of range **as tolerated**
 - Capsular stretches (especially posterior capsule if tight) **as tolerated**
- Continue to gradually restore pre-operative AAROM and then AROM with good SHR **as tolerated/not into pain**
- Gradually introduce and progress shoulder strengthening **as tolerated/not into pain**
 - Introduce scapular stabilisation exercises
 - Progress cuff strengthening, progressing from maximal isometrics to isotonic strengthening **as tolerated/not into pain**
 - Progress proprioceptive exercises
 - Gradually progress to shoulder and upper limb strengthening **as long as the patient is able to elevate the arm without the shoulder or scapular ‘hitching’**
 - Gradually progress to dynamic and rhythmic stabilisation exercises
- Progress lower limb and core strengthening as required

Precautions:

- Avoid excessive loading of shoulder
 - No heavy, repetitive or prolonged overhead activities

Milestones at 14 Weeks:

- Resolved pain, inflammation and muscle inhibition
- 100% of intra-operative, post-MUA/capsular release AROM with good SHR

Arthroscopic Capsular Release post-op Protocol		Classification: Company Confidential
Document type: Post-op Protocol	Page: 2 of 3	Author: Darren James
Effective Date: March 2022	Version: 2.1	Next Review: March 2024

Phase 4 (3 – 9 Months)

Goals:

- Maintain full PROM
 - Continue multi-directional stretching into end of range **as tolerated**
- Full pre-operative AROM with good SHR **as tolerated/not into pain**
- Progress shoulder strengthening **as tolerated/not into pain**
 - Progress cuff strengthening and scapular stabilisation exercises
 - Progress shoulder and upper limb strengthening ensuring good SHR
 - Progress dynamic and rhythmic stabilisation exercises
- Gradually progress functional activities
- Gradually progress sport-specific exercises

Milestones at 4 months:

- Full pain-free motion with good SHR

Sulis Specialist Orthopaedic Shoulder Team

Mr G Jennings (TR & Orth), Mr S Gregg-Smith (TR & Orth), Mr A Chamblor (TR & Orth) and Mr U Butt (Orth).

To contact the Consultants' Secretaries, call Sulis Hospital on 01761 422222 or to contact the Physiotherapy team call 01761 422388.

Arthroscopic Capsular Release post-op Protocol		Classification: Company Confidential
Document type: Post-op Protocol	Page: 3 of 3	Author: Darren James
Effective Date: March 2022	Version: 2.1	Next Review: March 2024